

Datasheet for ABIN966144

**anti-Fibulin 5 antibody****3** Publications[Go to Product page](#)

## Overview

|              |  |
|--------------|--|
| Quantity:    | 0.1 mL                                   |
| Target:      | Fibulin 5 (FBLN5)                        |
| Reactivity:  | Human                                    |
| Host:        | Please inquire                           |
| Clonality:   | Monoclonal                               |
| Conjugate:   | This Fibulin 5 antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA             |

## Product Details

|               |  |
|---------------|--|
| Isotype:      | IgM  |
| Specificity:  | Ni-NTA purified truncated recombinant Fibulin 5 expressed in E. Coli strain BL21 (DE3) |
| Purification: | Crude ascites.   |

## Target Details

|                   |   |
|-------------------|---|
| Target:           | Fibulin 5 (FBLN5)   |
| Alternative Name: | Fibulin 5 ( <a href="#">FBLN5 Products</a> )  |
| Background:       | Fibulin 5(FBLN5), with 448-amino acid protein (about 50kDa), is a recently discovered multifunctional extracellular matrix protein that mediates endothelial cell adhesion through integrin ligation, regulates cell growth and motility in a context-specific manner, and prevents elastinopathy in vivo. Fibulin-5 is abundantly expressed in great vessels and cardiac valves during embryogenesis, and in many adult tissues including the aorta, lung, uterus and skin, all of |

## Target Details

---

which contain abundant elastic fibres. Decreased fibulin-5 may contribute to the pathogenesis of aortic dissection by impairing elastic fiber assembly. Fibulin-5 is also a good marker of skin ageing and that the earlier loss of fibulin-5 may involve age-dependent changes in other elastic fibre components.

Pathways: [SARS-CoV-2 Protein Interactome](#)

## Application Details

---

Application Notes: Western Blot: 1: 500- 1: 2000  
ELISA: Propose dilution 1: 10,000.  
Determining optimal working dilutions by titration test.

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Storage: -20 °C

## Publications

---

Product cited in: Williams, Werner-Fraczek, Chang, Bailey-Serres: "Regulated phosphorylation of 40S ribosomal protein S6 in root tips of maize." in: **Plant physiology**, Vol. 132, Issue 4, pp. 2086-97, (2003) ([PubMed](#)).

McBride, Nemer: "The C-terminal domain of c-fos is required for activation of an AP-1 site specific for jun-fos heterodimers." in: **Molecular and cellular biology**, Vol. 18, Issue 9, pp. 5073-81, (1998) ([PubMed](#)).