# antibodies -online.com





# Datasheet for ABIN966145

# anti-Filamin-Like antibody



#### Overview

Quantity:	0.5 mL
Target:	Filamin-Like (LOC409697)
Reactivity:	Human, Chicken, Mouse, Rat
Host:	Mouse
Clonality:	Monoclonal
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Formalin-fixed Paraffin-embedded Sections) (IHC (fp))

#### **Product Details**

Immunogen:	Platelet Filamin
Clone:	FLMN01 (same as PM6-317)
Isotype:	lgG1
Purification:	Concentrated. Purified from the ascites fluid by Protein G chromatography

# Target Details

Target:	Filamin-Like (LOC409697)
Alternative Name:	Filamin (LOC409697 Products)
Background:	Filamin functions as a cross-linking protein forming a flexible link between two actin filaments. It is composed of two identical polypeptide chains each joined to the other at one end, with an actin binding site at the other. It is present in human platelets, lymphocytes, fibroblasts, and smooth muscle actin.

### **Target Details**

raiget Details	
	Also available without BSA and azide at 1mg/ml.
Molecular Weight:	250kDa
Application Details	
Application Notes:	ELISA (Order Ab without BSA)
	Western blotting: (Ab 1-2 micrograms/ml for 2hrs at RT)
	Immunohistology (Formalin/paraffin)(Use Ab at 1-2 micrograms/ml for 30 min at RT)
	Staining of formalin/paraffin tissues requires boiling tissue sections in 10mM citrate buffer, pH
	6.0, for 10-20 min followed by cooling at RT for 20 min.
Comment:	Cellular Localization: Cytoplasmic.
	Recommended Positive Control: NIH3T3 Cells, Skin
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.2 mg/mL.
Buffer:	10mM PBS, pH 7.4, with 0.2% BSA
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C