



[Go to Product page](#)

Datasheet for ABIN967283

## anti-Fc gamma RII (CD32) antibody

### 3 Images

#### Overview

Quantity:	0.5 mg
Target:	Fc gamma RII (CD32)
Reactivity:	Rat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	Un-conjugated
Application:	Flow Cytometry (FACS), Western Blotting (WB), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Blocking Reagent (BR), Immunohistochemistry (Zinc-fixed Sections) (IHC (zinc))

#### Product Details

Brand:	BD Pharmingen™
Immunogen:	Recombinant Rat CD32 Protein
Clone:	D34-485
Isotype:	IgG1 kappa
Characteristics:	<ol style="list-style-type: none"><li>1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.</li><li>2. Please refer to us for technical protocols.</li><li>3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.</li><li>4. Sodium azide is a reversible inhibitor of oxidative metabolism, therefore, antibody</li></ol>

## Product Details

---

preparations containing this preservative agent must not be used in cell cultures nor injected into animals. Sodium azide may be removed by washing stained cells or plate-bound antibody or dialyzing soluble antibody in sodium azide-free buffer. Since endotoxin may also affect the results of functional studies, we recommend the NA/LE™ (No Azide/Low Endotoxin) antibody format, if available, for in vitro and in vivo use.

**Purification:** The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

## Target Details

---

**Target:** Fc gamma RII (CD32)

**Alternative Name:** CD32 ([CD32 Products](#))

**Background:** The D34-485 antibody reacts specifically with CD32, the FcγRII receptor. Rat CD32 is expressed on B lymphocytes, myeloid cells, and some lymphocytes in the thymic medulla. D34-485 mAb blocks binding of aggregated immunoglobulins to the FcγRII receptors in vitro. This antibody is routinely tested by flow cytometric analysis.  
Synonyms: FcγRII Receptor

## Application Details

---

**Application Notes:** To specifically stain cells bearing Fc gamma II receptors for flow cytometric analysis: Incubate cell suspension with this antibody (Less or equal than 1 µg/million cells) followed by an appropriate fluorochrome-conjugated second-step reagent. To reduce Fc receptor-mediated binding by antibodies of interest to FcγRII receptor-bearing rat cells for flow cytometric analysis:

- Preincubate cell suspension with Rat Fc Block Solution (e.g., Less or equal than 1 µg/million cells in 100 µl) at 4°C for 5 minutes.
- Add antibody of interest directly to preincubated cells in the presence of Rat Fc Block Solution (i.e., Rat Fc Block Solution need not be washed off before staining cells).
- If anti-Ig second-step is necessary, a reagent must be chosen which will not bind to Rat Fc Block Solution (e.g., mouse IgG1, kappa)

**Restrictions:** For Research Use only

## Handling


---

**Format:** Liquid

## Handling


Concentration:	0.5 mg/mL
Buffer:	Aqueous buffered solution containing $\leq 0.09$ % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Store undiluted at 4° C.

## Images




### Flow Cytometry

**Image 1.** Blocking of Fc-mediated binding to Fcγ receptors (CD32) on rat splenocytes. Lewis rat splenocytes were pre-incubated with purified isotype control mAb A112-2 (first panel) or Rat BD Fc Block™ purified anti-rat CD32 mAb D34-485 (second panel). Two-color staining was performed with FITC-conjugated anti-rat NKR-P1A mAb 10/78 and PE-conjugated anti-rat CD45RA mAb OX-33. Note how the dim staining of B lymphocytes (OX-33+ cells) by anti-CD161a (NKR-P1A) (first panel) is reduced when Rat BD Fc Block™ is used before staining (second panel). Flow cytometry was performed on a BD FACScan™ flow cytometry system.



### Flow Cytometry

**Image 2.** Blocking of Fc-mediated binding to Fcγ receptors (CD32) on rat splenocytes. Lewis rat splenocytes were pre-incubated with purified isotype control mAb A112-2 (first panel) or Rat BD Fc Block™ purified anti-rat CD32 mAb D34-485 (second panel). Two-color staining was performed with FITC-conjugated anti-rat NKR-P1A mAb 10/78 and PE-conjugated anti-rat CD45RA mAb OX-33. Note how the dim staining of B lymphocytes (OX-33+ cells) by anti-CD161a (NKR-P1A) (first panel) is reduced when Rat BD



Fc Block™ is used before staining (second panel). Flow cytometry was performed on a BD FACScan™ flow cytometry system.

### Flow Cytometry

**Image 3.** Two-color analysis of the expression of CD32 on rat splenocytes. Lewis rat splenocytes were stained with purified D34-485 mAb, followed by FITC-conjugated polyclonal goat anti-mouse Ig, then PE-conjugated anti-rat IgM G53-238. Double-positive cells are B lymphocytes (IgM+ cells), which express CD32 on the cell surface. Flow cytometry was performed on a BD FACScan™ flow cytometry system.