antibodies -online.com





anti-rala antibody (AA 35-206)

4 Images



Publications



Go to Product page

_					
U	V	er	V	Ie	W

Quantity:	150 μg	
Target:	rala	
Binding Specificity:	AA 35-206	
Reactivity:	Rat, Dog, Chicken	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This rala antibody is un-conjugated	
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunoprecipitation (IP), Immunohistochemistry (Formalin-fixed Sections) (IHC (f))	

Product Details

Immunogen:	Human Ral A aa. 35-206	
Clone:	8-Ral A	
Isotype:	IgG2a kappa	
Cross-Reactivity:	Rat (Rattus), Chicken, Dog (Canine), Mouse (Murine)	
Characteristics:	1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.	
	2. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide	
	compounds in running water before discarding to avoid accumulation of potentially explosive	
	deposits in plumbing.	
	3. Source of all serum proteins is from USDA inspected abattoirs located in the United States.	
	4. Please refer to us for technical protocols.	

Product Details Purification: The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. **Target Details** Target: rala Alternative Name: Ral A (rala Products) Ral is a low molecular weight GTP-binding protein belonging to the Ras superfamily of GTP-Background: binding proteins and shows 50% amino acid identity to Ras. Ral cDNA clones have been isolated from human placenta, human pheochromocytoma, simian ß-lymphocyte, and marine ray electric lobe cDNA libraries. In humans, cDNA sequences for Ral A and RalB have been determined. The predicted amino acid sequences show 85% identity. Both proteins consist of 206 amino acids with a predicted molecular weight of 24kDa. However, Ral A isolated from human platelets shows an apparent molecular weight of 26-28kDa. Ral A mRNA levels in adult mouse tissues appear to be highest in testes, ovary, and brain, with lower levels found in the liver, spleen, kidney, thymus, heart, and salivary glands. Similar results have been observed in rat tissues. This antibody is routinely tested by western blot analysis. 24-28 kDa Molecular Weight: Pathways: Neurotrophin Signaling Pathway, SARS-CoV-2 Protein Interactome **Application Details** Comment: Related Products: ABIN968545, ABIN967389 Restrictions: For Research Use only Handling Format: Liquid Concentration: 250 µg/mL Buffer: Aqueous buffered solution containing BSA, glycerol, and ≤0.09 % sodium azide.

This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

Sodium azide

-20 °C

should be handled by trained staff only.

Preservative:

Storage:

Precaution of Use:

Storage Comment:

Store undiluted at -20°C.

Publications

Product cited in:

Xu, Frankel, Jackson, Rotunda, Boshans, DSouza-Schorey, Foster: "Elevated phospholipase D activity in H-Ras- but not K-Ras-transformed cells by the synergistic action of RalA and ARF6." in: **Molecular and cellular biology**, Vol. 23, Issue 2, pp. 645-54, (2003) (PubMed).


Kinashi, Katagiri, Watanabe, Vanhaesebroeck, Downward, Takatsu: "Distinct mechanisms of alpha 5beta 1 integrin activation by Ha-Ras and R-Ras." in: **The Journal of biological chemistry**, Vol. 275, Issue 29, pp. 22590-6, (2000) (PubMed).

Goi, Rusanescu, Urano, Feig: "Ral-specific guanine nucleotide exchange factor activity opposes other Ras effectors in PC12 cells by inhibiting neurite outgrowth." in: **Molecular and cellular biology**, Vol. 19, Issue 3, pp. 1731-41, (1999) (PubMed).

Polakis, Weber, Nevins, Didsbury, Evans, Snyderman: "Identification of the ral and rac1 gene products, low molecular mass GTP-binding proteins from human platelets." in: **The Journal of biological chemistry**, Vol. 264, Issue 28, pp. 16383-9, (1989) (PubMed).

Olofsson, Chardin, Touchot, Zahraoui, Tavitian: "Expression of the ras-related ralA, rho12 and rab genes in adult mouse tissues." in: **Oncogene**, Vol. 3, Issue 2, pp. 231-4, (1988) (PubMed).

Images




Immunofluorescence

Image 1. Immunofluorescence staining of rat neurons at 5 μ g/ml of the anti-Ral A antibody.

Image 2.





Western Blotting

Image 3.

Please check the product details page for more images. Overall 4 images are available for ABIN967836.