


# Datasheet for ABIN967926 anti-YES1 antibody (AA 10-193)





Go to Product page

$\sim$				
	Ive	r\/		٨
$\cup$	$\lor \lor \vdash$	I V I	$\Box$	٧V

Quantity:	50 μg	
Target:	YES1	
Binding Specificity:	AA 10-193	
Reactivity:	Human, Mouse, Rat, Dog, Chicken	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This YES1 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF)	
Product Details		
Immunogen:	Human Yes aa. 10-193	
Immunogen: Clone:	Human Yes aa. 10-193 1-Yes	
Clone:	1-Yes	
Clone: Isotype:	1-Yes IgG1	

chromatograph	١y.
---------------	-----

## Target Details

Target:	YES1	
Alternative Name:	Yes (YES1 Products)	
Background:	Yes is a member of the Src family of tyrosine kinases. The cellular gene was initially identified	
	as a homologue of v-yes, the oncogene of avian sarcoma virus Y73. At least nine different Src	
	family members have been identified. yes, fyn, and Src are widely expressed in a variety of cell	
	types, while the remaining members are primarily expressed in hematopoietic cells. Members	
	of this family have several common features: they have unique N-terminal domains, they attach	
	to cellular membranes through a myristylated N-terminus and they have homologus SH2, SH3,	
	and catalytic domains. The common expression patterns of Yes, fyn, and Src suggest some	
	overlapping functions. All three of these kinases can physically associate with a number of cell-	
	surface molecules which appear to increase their catalytic activity.	
Molecular Weight:	62 kDa	
Pathways:	CXCR4-mediated Signaling Events, Signaling Events mediated by VEGFR1 and VEGFR2,	
	Thromboxane A2 Receptor Signaling	

## **Application Details**

Comment:	Related Products: ABIN968533, ABIN967389
Restrictions:	For Research Use only
Handling	

панишну	
Format:	Liquid
Concentration:	250 μg/mL
Buffer:	Aqueous buffered solution containing BSA, glycerol, and ≤0.09 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store undiluted at -20°C.

Product cited in:

Chen, Lu, Goodenough, Jeansonne: "Nonreceptor tyrosine kinase c-Yes interacts with occludin during tight junction formation in canine kidney epithelial cells." in: **Molecular biology of the cell**, Vol. 13, Issue 4, pp. 1227-37, (2002) (PubMed).

Lee, Bonnah, Higashi, Atkinson, Milgram, So: "CD46 is phosphorylated at tyrosine 354 upon infection of epithelial cells by Neisseria gonorrhoeae." in: **The Journal of cell biology**, Vol. 156, Issue 6, pp. 951-7, (2002) (PubMed).

Encinas, Tansey, Tsui-Pierchala, Comella, Milbrandt, Johnson: "c-Src is required for glial cell line-derived neurotrophic factor (GDNF) family ligand-mediated neuronal survival via a phosphatidylinositol-3 kinase (PI-3K)-dependent pathway." in: **The Journal of neuroscience: the official journal of the Society for Neuroscience**, Vol. 21, Issue 5, pp. 1464-72, (2001) (PubMed).

Arizmendi, Family, Salas-Brito: "Quenched disorder effects on deterministic inertia ratchets." in: **Physical review. E, Statistical, nonlinear, and soft matter physics**, Vol. 63, Issue 6 Pt 1, pp. 61104, (2001) (PubMed).

Park, Cartwright: "Src activity increases and Yes activity decreases during mitosis of human colon carcinoma cells." in: **Molecular and cellular biology**, Vol. 15, Issue 5, pp. 2374-82, (1995) (PubMed).


There are more publications referencing this product on: Product page

#### **Images**



#### **Western Blotting**

**Image 1.** Western blot analysis of Yes on a A431 cell lysate (Human epithelial carcinoma, ATCC CRL-1555). Lane 1: 1:250, lane 2: 1:500, lane 3: 1:1000 dilution of the Mouse Anti-Yes antibody.



### **Immunohistochemistry**

**Image 2.** Immunohistochemical staining on a rabbit brain tissue section with the Mouse Anti-Yes antibody.