

Datasheet for ABIN967993

anti-SNAP91 antibody (AA 706-896)





Go to Product page

Overview

Quantity:	50 μg
Target:	SNAP91
Binding Specificity:	AA 706-896
Reactivity:	Human, Rat, Mouse
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This SNAP91 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunoprecipitation (IP)

Product Details

Immunogen:	Rat AP180 aa. 706-896
Clone:	34-AP180
Isotype:	lgG1
Cross-Reactivity:	Mouse (Murine), Human
Characteristics:	1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
	2. Please refer to us for technical protocols.
	3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide
	compounds in running water before discarding to avoid accumulation of potentially explosive
	deposits in plumbing.
	4. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

Product Details

			•				
ப	111	rı	+	icat	1/	71	\sim
	ш			ו ה. או	ш	. , ,	

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

Target Details

Target:	SNAP91
Alternative Name:	AP180 (SNAP91 Products)
Background:	AP180, also known as AP-3, F1-20, NP185, and pp155, is one of four assembly proteins which
	are involved in the organization and assembly of clathrin triskelia in clathrin-coated vesicles.
	Two of these proteins, AP-1 and AP-2, exist as tetramers, while AP180 and auxilin are
	monomeric assembly proteins. The clathrin binding potential of AP180 is defined by several
	regions of the molecule, including the 30 kDa N-terminal domain, a central domain, and a 58
	kDa C-terminal domain. Although clathrin binding occurs throughout the molecule, only the C-
	terminal domain is associated with both binding of clathrin and assembly of clathrin cages. The
	highly acidic central domain, which contains an uncharged alanine-rich segment, is thought to
	impart the irregular physical properties to this protein. AP180 is the only clathrin assembly
	protein specific for synapses and is thought to be involved in synaptic vesicle biogenesis and
	recycling. Furthermore, AP180 is bound by inositol-6-phosphate which has been shown to be
	closely regulated in neuronal cells by external stimuli.
	Synonyms: AP-3, F1-20, NP185, and pp155
Molecular Weight:	180 kDa

Application Details

Comment:

Storage:

Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	Aqueous buffered solution containing BSA, glycerol, and ≤0.09 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Precaution of Use:	·

Related Products: ABIN968545, ABIN967389

-20 °C

Storage Comment:

Store undiluted at -20° C.

Publications

Product cited in:

Kalthoff, Groos, Kohl, Mahrhold, Ungewickell: "Clint: a novel clathrin-binding ENTH-domain protein at the Golgi." in: **Molecular biology of the cell**, Vol. 13, Issue 11, pp. 4060-73, (2002) (PubMed).

Drake, Traub: "Interaction of two structurally distinct sequence types with the clathrin terminal domain beta-propeller." in: **The Journal of biological chemistry**, Vol. 276, Issue 31, pp. 28700-9, (2001) (PubMed).

Traub, Downs, Westrich, Fremont: "Crystal structure of the alpha appendage of AP-2 reveals a recruitment platform for clathrin-coat assembly." in: **Proceedings of the National Academy of Sciences of the United States of America**, Vol. 96, Issue 16, pp. 8907-12, (1999) (PubMed).

Ye, Lafer: "Clathrin binding and assembly activities of expressed domains of the synapse-specific clathrin assembly protein AP-3." in: **The Journal of biological chemistry**, Vol. 270, Issue 18, pp. 10933-9, (1995) (PubMed).


Morris, Schröder, Plessmann, Weber, Ungewickell: "Clathrin assembly protein AP180: primary structure, domain organization and identification of a clathrin binding site." in: **The EMBO journal**, Vol. 12, Issue 2, pp. 667-75, (1993) (PubMed).

Images



Western Blotting


Image 1. Western blot analysis of AP180 on a rat cerebrum lysate. Lane 1: 1:250, lane 2: 1:500, lane 3: 1:1000 dilution of the mouse anti-AP180 antibody.



Immunofluorescence

Image 2. Immunofluorescence staining of human endothelial cells.

Image 3.



Please check the product details page for more images. Overall 4 images are available for ABIN967993.