

Datasheet for ABIN968199

**anti-HAX1 antibody (AA 10-148)**

4 Images

1 Publication

[Go to Product page](#)

## Overview

Quantity:	150 µg
Target:	HAX1
Binding Specificity:	AA 10-148
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This HAX1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunoprecipitation (IP)

## Product Details

Immunogen:	Human HAX-1 aa. 10-148
Clone:	52-HAX
Isotype:	IgG1
Cross-Reactivity:	Rat (Rattus), Mouse (Murine)
Characteristics:	<ol style="list-style-type: none"><li>1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.</li><li>2. Please refer to us for technical protocols.</li><li>3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.</li><li>4. Source of all serum proteins is from USDA inspected abattoirs located in the United States.</li></ol>

## Product Details

Purification:	The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.
---------------	---

## Target Details

Target:	HAX1
Alternative Name:	HAX-1 ( <a href="#">HAX1 Products</a> )
Background:	Cell-cell interactions in the immune system, mediated by membrane-bound antigen receptors, trigger a cascade of signaling mechanisms collectively known as the T-cell response. Upon antigen binding, cellular tyrosine kinases are activated resulting in the phosphorylation of HS1 or Lck-binding protein 1 (LckBP1), which is involved in thymic negative selection. HS1 is one of the earliest intracellular proteins to be tyrosine phosphorylated in response to antigen-receptor cross-linking in T cells. HAX-1 (HS1-associated protein X-1) is a 35 kDa protein found associated with HS1 by co-immunoprecipitation assays and the two-hybrid system. HAX-1 is ubiquitously expressed and localized in the mitochondria, endoplasmic reticulum, and the nuclear envelope. The HS1-HAX-1 complex may be critical for signal transduction pathways initiated by antigen presentation to T cells.
Molecular Weight:	35 kDa
Pathways:	<a href="#">Regulation of Actin Filament Polymerization</a>

## Application Details

Comment:	Related Products: ABIN968537, ABIN967389
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	250 µg/mL
Buffer:	Aqueous buffered solution containing BSA, glycerol, and ≤0.09 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C

## Handling



Storage Comment: Store undiluted at -20° C.

## Publications

Product cited in: Suzuki, Demoliere, Kitamura, Takeshita, Deuschle, Watanabe: "HAX-1, a novel intracellular protein, localized on mitochondria, directly associates with HS1, a substrate of Src family tyrosine kinases." in: **Journal of immunology (Baltimore, Md. : 1950)**, Vol. 158, Issue 6, pp. 2736-44, (1997) ([PubMed](#)).


## Images

Image 1.



### Western Blotting

**Image 2.** Western blot analysis of HAX-1 on Jurkat cell lysate. Lane 1: 1:250, lane 2: 1:500, lane 3: 1:1000 dilution of anti-HAX-1 antibody.



#### Immunofluorescence

**Image 3.** Immunofluorescent staining of HeLa cells with anti-HAX-1 antibody.

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN968199.