

Datasheet for ABIN968250 anti-N-Cadherin antibody (AA 802-819)

4 Images

5 Publications



Overview

Quantity:	150 µg
Target:	N-Cadherin (CDH2)
Binding Specificity:	AA 802-819
Reactivity:	Human, Mouse, Rat, Chicken
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This N-Cadherin antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunoprecipitation (IP)

Product Details

Immunogen:	Mouse N-Cadherin aa. 802-819
Clone:	32-N
lsotype:	lgG1
Cross-Reactivity:	Human, Rat (Rattus), Chicken
Characteristics:	 Since applications vary, each investigator should titrate the reagent to obtain optimal results. Please refer to us for technical protocols. Source of all serum proteins is from USDA inspected abattoirs located in the United States. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
Purification:	The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN968250 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

Product Details

chromatography.

Target Details

Buffer:

Preservative:

Target:	N-Cadherin (CDH2)
Alternative Name:	N-Cadherin (CDH2 Products)
Background:	Cadherins are a family of Ca2+-dependent intercellular adhesion molecules that play a central
	role in controlling morphogenetic movements during development. Their function is regulated
	by association with the actin cytoskeleton by a complex of cytoplasmic proteins called the
	catenins (alpha, beta, gamma). Members of the cadherin family include P-cadherin, E-cadherin
	(uvomorulin), N-cadherin (neural cadherin), R-cadherin, cadherin 5, L-CAM, and EP-cadherin. N-
	cadherin mRNA is found at elevated levels in brain and heart and at a much lower level in liver.
	Mechanisms such as mRNA expression, cytokine modulation, and protease-mediated turnover
	modulate N-cadherin protein levels during development. In addition, N-cadherin function is
	indirectly regulated by endogenous kinases and phosphatases. Tyrosine phosphorylation of
	beta-catenin complexed with N-cadherin results in dissociation of N-cadherin from actin.
	However, N-cadherin also interacts with a PTP1B-like phosphatase that dephosphorylates beta-
	catenin and promotes N-cadherin/actin association. Thus, N-cadherin is an integral adhesion
	molecule whose function is regulated by protein-protein interactions and
	phosphorylation/dephosphorylation events.
Molecular Weight:	130 kDa
Pathways:	Regulation of Muscle Cell Differentiation, Cell-Cell Junction Organization, Synaptic Membrane
Application Details	
Comment:	Related Products: ABIN967389, ABIN968535
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	250 μg/mL

Aqueous buffered solution containing BSA, glycerol, and ≤ 0.09 % sodium azide.

Sodium azide

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN968250 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

Handling	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store undiluted at -20°C.
Publications	
Product cited in:	Izawa, Nishizawa, Ohtakara, Inagaki: "Densin-180 interacts with delta-catenin/neural plakophilin-
	related armadillo repeat protein at synapses." in: The Journal of biological chemistry, Vol. 277,
	Issue 7, pp. 5345-50, (2002) (PubMed).
	Nürnberger, Bacallao, Phillips: "Inversin forms a complex with catenins and N-cadherin in
	polarized epithelial cells." in: Molecular biology of the cell, Vol. 13, Issue 9, pp. 3096-106, (2002)
	(PubMed).
	Bhowmick, Ghiassi, Bakin, Aakre, Lundquist, Engel, Arteaga, Moses: "Transforming growth
	factor-beta1 mediates epithelial to mesenchymal transdifferentiation through a RhoA-
	dependent mechanism." in: Molecular biology of the cell, Vol. 12, Issue 1, pp. 27-36, (2001) (
	PubMed).
	Cowin: "Unraveling the cytoplasmic interactions of the cadherin superfamily." in: Proceedings
	of the National Academy of Sciences of the United States of America, Vol. 91, Issue 23, pp.
	10759-61, (1994) (PubMed).
	Miyatani, Shimamura, Hatta, Nagafuchi, Nose, Matsunaga, Hatta, Takeichi: "Neural cadherin:
	role in selective cell-cell adhesion." in: Science (New York, N.Y.), Vol. 245, Issue 4918, pp. 631-5,
	(1989) (PubMed).





Western Blotting

Image 1. Western blot analysis of N-Cadherin on HeLa lysate. Lane 1: 1:250, lane 2: 1:500, lane 3: 1:1000, lane 4: 1:2000 dilution of anti-N-Cadherin.

Immunofluorescence

Image 2. Immunofluorescent staining of 293 cells with anti-N-Cadherin.

Image 3.

Generated from human N-Cadherin



Please check the product details page for more images. Overall 4 images are available for ABIN968250.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 4/4 | Product datasheet for ABIN968250 | 07/26/2024 | Copyright antibodies-online. All rights reserved.