antibodies - online.com





anti-PRKACA antibody (AA 18-347)

3 Images



Publications



Go to Product page

| () | 11/0 | r\ /1 | $\triangle 1 $ |
|-----|--------------------|------------|----------------|
| | $\lor \lor \vdash$ | $I \vee I$ | ew |
| | | | |

| Quantity: | 150 μg |
|----------------------|--|
| Target: | PRKACA |
| Binding Specificity: | AA 18-347 |
| Reactivity: | Human, Mouse, Rat, Dog |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Conjugate: | This PRKACA antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunofluorescence (IF) |

Product Details

| Immunogen: | Human PKA[Calpha] subunit aa. 18-347 | |
|-------------------|---|--|
| Clone: | 5B | |
| Isotype: | lgG2b | |
| Cross-Reactivity: | Dog (Canine), Rat (Rattus), Mouse (Murine) | |
| Characteristics: | Since applications vary, each investigator should titrate the reagent to obtain optimal results. Please refer to us for technical protocols. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing. | |
| | 4. Source of all serum proteins is from USDA inspected abattoirs located in the United States. | |
| Purification: | The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity | |

chromatography.

Target Details

| Target: | PRKACA | |
|---------------------|---|--|
| Alternative Name: | PKA C (PRKACA Products) | |
| Background: | CAMP-dependent Protein Kinase (PKA) is composed of two distinct subunits: catalytic (C) and regulatory (R). Four regulatory subunits have been identified: RIalpha, RIbeta, RIIalpha, and RIIbeta. These subunits define type I and II cAMP-dependent protein kinases. Following binding of cAMP, the regulatory subunits dissociate from the catalytic subunits, rendering the enzyme active. Type I and type II holoenzymes have three potential C subunits (Calpha, Cbeta, or | |
| Molecular Weight: | Cgamma). Type II PKA can be distinguished by autophosphorylation of the R-subunits, while type I PKA binds Mg/ATP with high affinity. The levels of expression of the different subunits vary according to cell and tissue type. 40 kDa | |
| Pathways: | NF-kappaB Signaling, Hedgehog Signaling, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Thyroid Hormone Synthesis, Carbohydrate Homeostasis, Myometrial Relaxation and Contraction, M Phase, G-protein mediated Events, Signaling Events mediated by VEGFR1 and VEGFR2, Interaction of EGFR with phospholipase C-gamma, Thromboxane A2 Receptor Signaling, VEGFR1 Specific Signals, Lipid Metabolism, SARS-CoV-2 Protein Interactome, The Global Phosphorylation Landscape of SARS-CoV-2 Infection | |
| Application Details | | |

Application Details

Comment:

Preservative:

Precaution of Use:

| Restrictions: | For Research Use only | |
|----------------|---|--|
| Handling | | |
| Format: | Liquid | |
| Concentration: | 250 μg/mL | |
| Buffer: | Aqueous buffered solution containing BSA, glycerol, and ≤0.09 % sodium azide. | |
| | | |

Related Products: ABIN968535, ABIN967389

Sodium azide

This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

Handling

| | should be handled by trained staff only. |
|------------------|--|
| Storage: | -20 °C |
| Storage Comment: | Store undiluted at -20°C. |
| Publications | |

Product cited in:


DeSouza, Reiken, Ondrias, Yang, Matkovich, Marks: "Protein kinase A and two phosphatases are components of the inositol 1,4,5-trisphosphate receptor macromolecular signaling complex." in: **The Journal of biological chemistry**, Vol. 277, Issue 42, pp. 39397-400, (2002) (PubMed).

Chen, Yu, Lee, Chiang, Chao, Huang, Chiong, Huang, Lai, Yang-Yen, Yen: "CREB is one component of the binding complex of the Ces-2/E2A-HLF binding element and is an integral part of the interleukin-3 survival signal." in: **Molecular and cellular biology**, Vol. 21, Issue 14, pp. 4636-46, (2001) (PubMed).

Orellana, Marfella-Scivittaro: "Distinctive cyclic AMP-dependent protein kinase subunit localization is associated with cyst formation and loss of tubulogenic capacity in Madin-Darby canine kidney cell clones." in: **The Journal of biological chemistry**, Vol. 275, Issue 28, pp. 21233-40, (2000) (PubMed).


Westphal, Soderling, Alto, Langeberg, Scott: "Scar/WAVE-1, a Wiskott-Aldrich syndrome protein, assembles an actin-associated multi-kinase scaffold." in: **The EMBO journal**, Vol. 19, Issue 17, pp. 4589-600, (2000) (PubMed).

Taylor, Buechler, Yonemoto: "cAMP-dependent protein kinase: framework for a diverse family of regulatory enzymes." in: **Annual review of biochemistry**, Vol. 59, pp. 971-1005, (1990) (PubMed).



Western Blotting

Image 1. Western blot analysis of PKA[C] on HeLa cell lysate. Lane 1: 1:1000, lane 2: 1:2000, lane 3: 1:4000 dilution of antiPKA[C].



Immunofluorescence

Image 2. Immunofluorescent staining of HeLa cells.

Image 3.

