



[Go to Product page](#)

Datasheet for ABIN968971
anti-Aurora A antibody

4 Images

2 Publications

Overview

Quantity:	100 µL
Target:	Aurora A (AURKA)
Reactivity:	Human, Rat, Monkey
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Aurora A antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Flow Cytometry (FACS)

Product Details

Immunogen:	Purified recombinant fragment of human AURKA expressed in E. coli.
Clone:	1F8
Isotype:	IgG1
Purification:	purified

Target Details

Target:	Aurora A (AURKA)
Alternative Name:	AURKA (AURKA Products)
Background:	Description: Aurora A plays a role in cell cycle regulation during anaphase and/or telophase, in relation to the function of the centrosome/spindle pole region during chromosome segregation. Aurora A plays a key role during tumor development and progression and is overexpressed in many human cancers including breast, ovarian and colorectal. Aurora A is viewed as a potential

Target Details

target for anticancer drug treatment. Tissue specificity: Highly expressed in testis and weakly in skeletal muscle, thymus and spleen. Also highly expressed in colon, ovarian, prostate, neuroblastoma, breast and cervical cancer cell lines.

Aliases: AIK, ARK1, AURA, BTAK, STK6, STK7, STK15, AURORA2, MGC34538, AURKA

Molecular Weight: 48 kDa

Gene ID: 6790

HGNC: 6790

Pathways: [Cell Division Cycle](#), [Asymmetric Protein Localization](#)

Application Details

Application Notes: ELISA: 1:10000, WB: 1:500 - 1:2000, ICC: 1:200 - 1:1000, FCM: 1:200 - 1:400

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Ascitic fluid containing 0.03 % sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C/-20 °C


Storage Comment: 4°C, -20°C for long term storage

Publications

Product cited in: Sproul, Xu, Wilhelm, Gire, Greene: "Cbl negatively regulates JNK activation and cell death." in: **Cell research**, Vol. 19, Issue 8, pp. 950-61, (2009) ([PubMed](#)).


Sanada, Suzuki, Shih, Otsu, Kato, Yamazaki, Tamura, Honda, Sakata-Yanagimoto, Kumano, Oda, Yamagata, Takita, Gotoh, Nakazaki, Kawamata, Onodera, Nobuyoshi, Hayashi, Harada, Kurokawa, Chiba, Mori et al.: "Gain-of-function of mutated C-CBL tumour suppressor in myeloid neoplasms. ..." in: **Nature**, Vol. 460, Issue 7257, pp. 904-8, (2009) ([PubMed](#)).

Loh, Sakai, Flotho, Kang, Fliegau, Archambeault, Mullighan, Chen, Bergstraesser, Bueso-Ramos, Emanuel, Hasle, Issa, van den Heuvel-Eibrink, Locatelli, Stary, Trebo, Wlodarski, Zecca, Shannon et al.: "Mutations in CBL occur frequently in juvenile myelomonocytic leukemia. ..." in: **Blood**, Vol. 114, Issue 9, pp. 1859-63, (2009) ([PubMed](#)).




ELISA

Image 1. Red: Control Antigen (100 ng), Purple: Antigen (10 ng), Green: Antigen (50 ng), Blue: Antigen (100 ng),



Flow Cytometry

Image 2. Flow cytometric analysis of K562 cells using AURKA mouse mAb (green) and negative control (purple).



Western Blotting

Image 3. Western blot analysis using AURKA mouse mAb against HEK293 (1), Sw620 (2), MCF-7 (3), Jurkat (4), Hela (5), HepG2 (6), Cos7 (7) and PC-12 (8) cell lysate.

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN968971.