antibodies -online.com







anti-TNFRSF8 antibody

Images



Publication



Overview

Quantity:	100 μL
Target:	TNFRSF8
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This TNFRSF8 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Western Blotting (WB), ELISA

Product Details

Immunogen:	Purified recombinant fragment of human CD30 expressed in E. coli.
Clone:	3B10
Isotype:	lgG1
Purification:	purified

Target Details

Target:	TNFRSF8
Alternative Name:	CD30 (TNFRSF8 Products)
Background:	Description: The protein encoded by this gene is a member of the TNF-receptor superfamily.
	This receptor is expressed by activated, but not by resting, T and B cells. TRAF2 and TRAF5 can
	interact with this receptor, and mediate the signal transduction that leads to the activation of
	NF-kappaB. This receptor is a positive regulator of apoptosis, and also has been shown to limit

Target Details

the proliferative potential of autoreactive CD8 effector T cells and protect the body against	
autoimmunity. Two alternatively spliced transcript variants of this gene encoding distinct	
isoforms have been reported.	
Aliases: CD30, Ki-1, D1S166E, TNFRSF8	

Gene ID: 943

HGNC: 943

Application Details

Application Notes:	ELISA: 1:10000, WB: 1:500 - 1:2000, FCM: 1:200 - 1:400

Restrictions: For Research Use only

Handling

Format:	Liquid
Buffer:	Ascitic fluid containing 0.03 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C/-20 °C
Storage Comment:	4°C, -20°C for long term storage


Publications

Product cited in:

Mishra, Thakur, Somal, Parmar, Yadav, Bharati, Bharti, Paul, Verma, Chouhan, Sharma, Singh, González, DOcchio, Sarkar et al.: "Expression and localization of angiopoietin family in buffalo ovarian follicles during different stages of development and modulatory role of angiopoietins on steroidogenesis and survival of cultured ..." in: **Theriogenology**, Vol. 86, Issue 7, pp. 1818-33, (2016) (PubMed).


Mishra, Parmar, Yadav, Reshma, Bharati, Bharti, Paul, Chouhan, Taru Sharma, Singh, Sarkar et al.: "Expression and localization of angiopoietin family in corpus luteum during different stages of oestrous cycle and modulatory role of angiopoietins on steroidogenesis, angiogenesis and survivability ..." in: **Reproduction in domestic animals = Zuchthygiene**, Vol. 51, Issue 6, pp. 855-869, (2016) (PubMed).

Images




ELISA

Image 1. Black line: Control Antigen (100 ng), Purple line: Antigen(10 ng), Blue line: Antigen (50 ng), Red line: Antigen (100 ng),



Flow Cytometry

Image 2. Flow cytometric analysis of Hela cells using CD30 mouse mAb (green) and negative control (red).



Western Blotting

Image 3. Western blot analysis using CD30 mouse mAb against Hela cell lysate.

Please check the product details page for more images. Overall 4 images are available for ABIN969019.