

Datasheet for ABIN969087
anti-Dynamin 1 antibody[Go to Product page](#)

2 Images

2 Publications

Overview

Quantity:	100 µL
Target:	Dynamin 1 (DNM1)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Dynamin 1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	Purified recombinant fragment of human Dynamin-1 expressed in E. coli.
Clone:	3G4B6
Isotype:	IgG2a
Purification:	purified

Target Details

Target:	Dynamin 1 (DNM1)
Alternative Name:	Dynamin-1 (DNM1 Products)
Background:	<p>Description: Dynamin-1 (Dyn1), with 864-amino acid protein (about 95 kDa), belongs to the dynamin family. Dynamin-1 (neuron-specific), dynamin-2 (ubiquitously expressed), and dynamin-3 (expressed only in the testis, brain, and lung), constitute the dynamin family.</p> <p>Members of the dynamin family are GTPase, microtubule-associated proteins which are</p>

Target Details

involved in endocytosis, synaptic transmission and neurogenesis. Dynamin-1 is phosphorylated in nerve terminals exclusively in the cytosolic compartment and in vitro by protein kinase C. Dynamin-1 is a large GTPase enzyme required in membrane constriction and fission during multiple forms of endocytosis. Dynamin-1 is also a key molecule required for the recycling of synaptic vesicles in neurons, and it has been known that dynamin-1 gene expression is induced during neuronal differentiation.

Aliases: DNM, DNM1

Molecular Weight: 97 kDa

Gene ID: 1759

HGNC: 1759

Pathways: [Toll-Like Receptors Cascades](#), [CXCR4-mediated Signaling Events](#), [Thromboxane A2 Receptor Signaling](#)

Application Details

Application Notes: ELISA: 1:10000, WB: 1:500 - 1:2000, IHC: 1:200 - 1:1000

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Ascitic fluid containing 0.03 % sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C/-20 °C

Storage Comment: 4°C, -20°C for long term storage

Publications

Product cited in: Gertych, Oh, Wawrowsky, Weisenberger, Tajbakhsh: "3-D DNA methylation phenotypes correlate with cytotoxicity levels in prostate and liver cancer cell models." in: **BMC pharmacology & toxicology**, Vol. 14, pp. 11, (2013) ([PubMed](#)).


Tajbakhsh: "Covisualization of methylcytosine, global DNA, and protein biomarkers for In Situ 3D DNA methylation phenotyping of stem cells." in: **Methods in molecular biology (Clifton, N.J.)**, Vol. 1052, pp. 77-88, (2013) ([PubMed](#)).

Fukuda, Ichiyanagi, Yamada, Go, Udono, Wada, Maeda, Soejima, Saitou, Ito, Sasaki: "Regional DNA methylation differences between humans and chimpanzees are associated with genetic changes, transcriptional divergence and disease genes." in: **Journal of human genetics**, Vol. 58, Issue 7, pp. 446-54, (2013) ([PubMed](#)).

Kurita, Arai, Nakamoto, Kato, Niwa: "Determination of DNA methylation using electrochemiluminescence with surface accumulable coreactant." in: **Analytical chemistry**, Vol. 84, Issue 4, pp. 1799-803, (2012) ([PubMed](#)).


Kurita, Niwa: "DNA methylation analysis triggered by bulge specific immuno-recognition." in: **Analytical chemistry**, Vol. 84, Issue 17, pp. 7533-8, (2012) ([PubMed](#)).

Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of paraffin-embedded human lymph tissue (A), glioma tissue (B) and cerebellum tissue (C), showing membrane localization using Dynamin1 mouse mAb with DAB staining



Western Blotting

Image 2. Western blot analysis using Dynamin1 mouse mAb against C6 (1), NIH/3T3 (2), SKN-SH (3), LN18 (4), SHSY5Y (5) cell lysate and rat brain tissues lysate (6).