antibodies -online.com







anti-FGG antibody



Publications



0	1 /	-	K	/1	-	1 A
u	\/	\vdash	ı ۱	/ I	\vdash	1/1

Quantity:	100 μL
Target:	FGG
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This FGG antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA, Immunocytochemistry (ICC), Flow Cytometry (FACS)

Product Details

Immunogen:	Purified recombinant fragment of human FGG expressed in E. coli.
Clone:	4H9
Isotype:	IgG2a
Purification:	purified

Target Details

Target:	FGG
Alternative Name:	FGG (FGG Products)
Background:	Description: The protein encoded by this gene is the gamma component of fibrinogen, a blood- borne glycoprotein comprised of three pairs of nonidentical polypeptide chains. Following vascular injury, fibrinogen is cleaved by thrombin to form fibrin which is the most abundant

component of blood clots. In addition, various cleavage products of fibrinogen and fibrin
regulate cell adhesion and spreading, display vasoconstrictor and chemotactic activities, and
are mitogens for several cell types. Mutations in this gene lead to several disorders, including
dysfibrinogenemia, hypofibrinogenemia and thrombophilia. Alternative splicing results in two
transcript variants encoding different isoforms.

Aliases: FGG

Molecular Weight:	52 kDa
Gene ID:	2266
HGNC:	2266

Application Details

Application Notes:	ELISA: 1:10000, WB: 1:500 - 1:2000, IHC: 1:200 - 1:1000, ICC: 1:200 - 1:1000, FCM: 1:200 - 1:400
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Ascitic fluid containing 0.03 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C/-20 °C
Storage Comment:	4°C, -20°C for long term storage

Publications

Product cited in:

Gertych, Oh, Wawrowsky, Weisenberger, Tajbakhsh: "3-D DNA methylation phenotypes correlate with cytotoxicity levels in prostate and liver cancer cell models." in: **BMC pharmacology & toxicology**, Vol. 14, pp. 11, (2013) (PubMed).


Tajbakhsh: "Covisualization of methylcytosine, global DNA, and protein biomarkers for In Situ 3D DNA methylation phenotyping of stem cells." in: **Methods in molecular biology (Clifton, N.J.)**, Vol. 1052, pp. 77-88, (2013) (PubMed).

Fukuda, Ichiyanagi, Yamada, Go, Udono, Wada, Maeda, Soejima, Saitou, Ito, Sasaki: "Regional DNA methylation differences between humans and chimpanzees are associated with genetic changes, transcriptional divergence and disease genes." in: **Journal of human genetics**, Vol. 58, Issue 7, pp. 446-54, (2013) (PubMed).

Kurita, Arai, Nakamoto, Kato, Niwa: "Determination of DNA methylation using electrochemiluminescence with surface accumulable coreactant." in: **Analytical chemistry**, Vol. 84, Issue 4, pp. 1799-803, (2012) (PubMed).


Kurita, Niwa: "DNA methylation analysis triggered by bulge specific immuno-recognition." in: **Analytical chemistry**, Vol. 84, Issue 17, pp. 7533-8, (2012) (PubMed).

Images




ELISA

Image 1. Red: Control Antigen (100 ng), Purple: Antigen (10 ng), Green: Antigen (50 ng), Blue: Antigen (100 ng),



Flow Cytometry

Image 2. Flow cytometric analysis of HepG2 cells using FGG mouse mAb (blue) and negative control (red).



Immunohistochemistry

Image 3. Immunohistochemical analysis of paraffinembedded cerebellum tissues using FGG mouse mAb with DAB staining.

Please check the product details page for more images. Overall 6 images are available for ABIN969142.