

Datasheet for ABIN969441

anti-TNNI2 antibody

7 Images

2 Publications

[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	TNNI2
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This TNNI2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Flow Cytometry (FACS)

Product Details

Immunogen:	Purified recombinant fragment of human TNNI2 expressed in E. coli.
Clone:	2F12A11
Isotype:	IgG1
Purification:	purified

Target Details

Target:	TNNI2
Alternative Name:	TNNI2 (TNNI2 Products)
Background:	Description: This gene encodes a fast-twitch skeletal muscle protein, a member of the troponin I gene family, and a component of the troponin complex including troponin T, troponin C and troponin I subunits. The troponin complex, along with tropomyosin, is responsible for the calcium-dependent regulation of striated muscle contraction. Mouse studies show that this

Target Details

component is also present in vascular smooth muscle and may play a role in regulation of smooth muscle function. In addition to muscle tissues, this protein is found in corneal epithelium, cartilage where it is an inhibitor of angiogenesis to inhibit tumor growth and metastasis, and mammary gland where it functions as a co-activator of estrogen receptor-related receptor alpha. This protein also suppresses tumor growth in human ovarian carcinoma. Mutations in this gene cause myopathy and distal arthrogryposis type 2B. Alternatively spliced transcript variants have been found for this gene.

Aliases: DA2B, FSSV, fsTnl, AMCD2B

Molecular Weight: 21 kDa

Gene ID: 7136

HGNC: 7136

Application Details

Application Notes: ELISA: 1:10000, WB: 1:500 - 1:2000, IHC: 1:200 - 1:1000, FCM: 1:200 - 1:400

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Ascitic fluid containing 0.03 % sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C/-20 °C

Storage Comment: 4°C, -20°C for long term storage

Publications

Product cited in: Gertych, Oh, Wawrowsky, Weisenberger, Tajbakhsh: "3-D DNA methylation phenotypes correlate with cytotoxicity levels in prostate and liver cancer cell models." in: **BMC pharmacology & toxicology**, Vol. 14, pp. 11, (2013) ([PubMed](#)).


Tajbakhsh: "Covisualization of methylcytosine, global DNA, and protein biomarkers for In Situ

3D DNA methylation phenotyping of stem cells." in: **Methods in molecular biology (Clifton, N.J.)**, Vol. 1052, pp. 77-88, (2013) ([PubMed](#)).

Fukuda, Ichiyangi, Yamada, Go, Udono, Wada, Maeda, Soejima, Saitou, Ito, Sasaki: "Regional DNA methylation differences between humans and chimpanzees are associated with genetic changes, transcriptional divergence and disease genes." in: **Journal of human genetics**, Vol. 58, Issue 7, pp. 446-54, (2013) ([PubMed](#)).


Kurita, Arai, Nakamoto, Kato, Niwa: "Determination of DNA methylation using electrochemiluminescence with surface accumulable coreactant." in: **Analytical chemistry**, Vol. 84, Issue 4, pp. 1799-803, (2012) ([PubMed](#)).

Kurita, Niwa: "DNA methylation analysis triggered by bulge specific immuno-recognition." in: **Analytical chemistry**, Vol. 84, Issue 17, pp. 7533-8, (2012) ([PubMed](#)).




ELISA

Image 1. Red: Control Antigen (100 ng), Purple: Antigen (10 ng), Green: Antigen (50 ng), Blue: Antigen (100 ng),



Flow Cytometry

Image 2. Flow cytometric analysis of NIH/3T3 cells using TNNI2 mouse mAb (blue) and negative control (red).



Immunohistochemistry

Image 3. Immunohistochemical analysis of paraffin-embedded rabbit cardiac muscle tissues using TNNI2 mouse mAb with DAB staining.

Please check the [product details page](#) for more images. Overall 7 images are available for ABIN969441.