



[Go to Product page](#)

Datasheet for ABIN969456

anti-WDR5 antibody

1 Image

2 Publications

Overview

Quantity:	100 µL
Target:	WDR5
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Application:	Western Blotting (WB), ELISA

Product Details

Immunogen:	Purified recombinant fragment of human WDR5 expressed in E. coli.
Clone:	7B11
Isotype:	IgG2b
Purification:	purified

Target Details

Target:	WDR5
Alternative Name:	WDR5 (WDR5 Products)
Background:	Description: WD-repeat protein 5 (WDR5, also designated BMP-2-induced gene 3 kb or BIG-3) belongs to the family of WD-40 repeat proteins, and is essential for vertebrate development, Hox gene activation and global H3K4 trimethylation. WDR5 is a conserved subunit of Trithorax (TRX) histone methyltransferase complexes that selectively binds to dimethylated Lys4 (K4me2) in Histone H3 to promote K4 trimethylation by TRX. It is expressed in osteoblasts,

Target Details

chondrocytes, osteocytes and marrow stromal cells. The WDR5 protein contains 7 WD-repeats, which may play a role in its function of accelerating osteoblast differentiation.

Aliases: SWD3, BIG-3, WDR5

Molecular Weight: 36.6 kDa

Gene ID: 11091

HGNC: 11091

Application Details

Application Notes: ELISA: 1:10000, WB: 1:500 - 1:2000

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Ascitic fluid containing 0.03 % sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: 4 °C/-20 °C

Storage Comment: 4°C, -20°C for long term storage

Publications

Product cited in: Jan, Adolfsson, Allaman, Buccarello, Magistretti, Pfeifer, Muhs, Lashuel: "Abeta42 neurotoxicity is mediated by ongoing nucleated polymerization process rather than by discrete Abeta42 species." in: **The Journal of biological chemistry**, Vol. 286, Issue 10, pp. 8585-96, (2011) ([PubMed](#)).

Deshmukh, Salehzadeh, Metayer-Coustard, Fahlman, Nair, Al-Khalili: "Post-transcriptional gene silencing of ribosomal protein S6 kinase 1 restores insulin action in leucine-treated skeletal muscle." in: **Cellular and molecular life sciences : CMLS**, Vol. 66, Issue 8, pp. 1457-66, (2009) ([PubMed](#)).



Western Blotting

Image 1. Western blot analysis using WDR5 mouse mAb against HeLa (1) cell lysate.