



[Go to Product page](#)

Datasheet for ABIN969526

anti-Hexosaminidase A antibody

4 Images

2 Publications

Overview

Quantity:	0.1 mg
Target:	Hexosaminidase A (HEXA)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Hexosaminidase A antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Flow Cytometry (FACS)

Product Details

Immunogen:	Purified recombinant fragment of human HEXA expressed in E. coli.
Clone:	3F10
Isotype:	IgG2b
Purification:	purified

Target Details

Target:	Hexosaminidase A (HEXA)
Alternative Name:	HEXA (HEXA Products)
Background:	Description: This gene encodes the alpha subunit of the lysosomal enzyme beta-hexosaminidase that, together with the cofactor GM2 activator protein, catalyzes the degradation of the ganglioside GM2, and other molecules containing terminal N-acetylhexosamines. Beta-hexosaminidase is composed of two subunits, alpha and beta, which are

Target Details

encoded by separate genes. Both beta-hexosaminidase alpha and beta subunits are members of family 20 of glycosyl hydrolases. Mutations in the alpha or beta subunit genes lead to an accumulation of GM2 ganglioside in neurons and neurodegenerative disorders termed the GM2 gangliosidoses. Alpha subunit gene mutations lead to Tay-Sachs disease (GM2-gangliosidosis type I).

Aliases: TSD

Molecular Weight: 60.7 kDa

Gene ID: 3073

HGNC: 3073

Pathways: [Sensory Perception of Sound, Glycosaminoglycan Metabolic Process](#)

Application Details

Application Notes: ELISA: 1:10000, WB: 1:500 - 1:2000, FCM: 1:200 - 1:400

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Purified antibody in PBS with 0.05 % sodium azide

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C/-20 °C

Storage Comment: 4°C, -20°C for long term storage

Publications

Product cited in: Gertych, Oh, Wawrowsky, Weisenberger, Tajbakhsh: "3-D DNA methylation phenotypes correlate with cytotoxicity levels in prostate and liver cancer cell models." in: **BMC pharmacology & toxicology**, Vol. 14, pp. 11, (2013) ([PubMed](#)).

Tajbakhsh: "Covisualization of methylcytosine, global DNA, and protein biomarkers for In Situ 3D DNA methylation phenotyping of stem cells." in: **Methods in molecular biology (Clifton,**


N.J., Vol. 1052, pp. 77-88, (2013) ([PubMed](#)).

Fukuda, Ichiyonagi, Yamada, Go, Udono, Wada, Maeda, Soejima, Saitou, Ito, Sasaki: "Regional DNA methylation differences between humans and chimpanzees are associated with genetic changes, transcriptional divergence and disease genes." in: **Journal of human genetics**, Vol. 58, Issue 7, pp. 446-54, (2013) ([PubMed](#)).

Kurita, Arai, Nakamoto, Kato, Niwa: "Determination of DNA methylation using electrochemiluminescence with surface accumulative coreactant." in: **Analytical chemistry**, Vol. 84, Issue 4, pp. 1799-803, (2012) ([PubMed](#)).


Kurita, Niwa: "DNA methylation analysis triggered by bulge specific immuno-recognition." in: **Analytical chemistry**, Vol. 84, Issue 17, pp. 7533-8, (2012) ([PubMed](#)).

Images




ELISA

Image 1. Black line: Control Antigen (100 ng), Purple line: Antigen(10 ng), Blue line: Antigen (50 ng), Red line: Antigen (100 ng),



Flow Cytometry

Image 2. Flow cytometric analysis of HeLa cells using HEXA mouse mAb (green) and negative control (red).



Western Blotting

Image 3. Western blot analysis using HEXA mouse mAb against L1210 (1), and HL7702 (2) cell lysate.

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN969526.