

Datasheet for ABIN969886 Rabbit anti-Human IgM (Chain mu), (Heavy Chain) Antibody (HRP)



Overview Quantity: 1 mg Target: ΙgΜ Binding Specificity: Chain mu, Heavy Chain Reactivity: Human Rabbit Host: HRP Conjugate: Application: ELISA, Immunohistochemistry (IHC), Western Blotting (WB) **Product Details** Purified Human IgM (µ chain) Immunogen: Based on IEP, this antibody reacts with: heavy (μ) chains on human IgM Specificity: No Cross-Reactivity: Human

| Characteristics: | 1.50 mg/mL (E 1 % at 280 nm = 13.0) |
|------------------|-------------------------------------|
| Purity: | > 95 % based on SDS-PAGE |

Target Details

| Target: | lgM |
|--------------|--------------|
| Abstract: | IgM Products |
| Target Type: | Antibody |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN969886 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

| Application Details | |
|---------------------|--|
| Application Notes: | This conjugate is suitable for all immunoassay applications. The optimal working dilution should be determined by the investigator. Suggested starting dilution: 1:500-1:5,000 for Immunohistochemistry. 1:200-1:5000 for ELISA/Western blot |
| Comment: | Country of Origin: Rabbit serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Lyophilized |
| Reconstitution: | Rehydrate with 1.1 mL of deionized water. Allow reconstituted product to stand for at least 30 minutes at room temperature prior to dilution. If necessary, Centrifuge to remove any particulates. Prepare fresh working dilution daily. |
| Concentration: | 1.0 mg/mL |
| Buffer: | 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free |
| Preservative: | Kathon CG |
| Handling Advice: | Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase. |
| Storage: | 4 °C |