# antibodies .- online.com







## **Normal Syrian Hamster Serum**



| $\sim$ |     |     |     |
|--------|-----|-----|-----|
|        | N/P | r\/ | i⊢₩ |

| Quantity:    | 2 mL   |  |
|--------------|--|--|
| Host:        | Hamster  |  |
| Application: | ELISA, Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Flow Cytometry (FACS), Labeling (LbI), Blocking Reagent (BR), Control (Ct) |  |

#### **Product Details**

| Specificity:     | Non-immune  |  |
|------------------|---|--|
| Characteristics: | Concentration: 50.0 mg/ml (Bradford, IgG standards)   |  |
| Purification:    | Lipid extracted and dialyzed against 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |  |

### **Application Details**

| Application Notes: | This product is used as a blocking reagent or control for most immunoassay applications. |
|--------------------|--|
| Restrictions:      | For Research Use only  |

#### Handling

| Format:         | Liquid   |  |
|-----------------|--|--|
| Reconstitution: | Rehydrate with 2.0 mL of deionized water. Swirl gentle and let stand for up to 2 hours at 18-25 °C. Centrifuge reconstituted serum to remove any precipitates. |  |
| Concentration:  | 50.0 mg/mL   |  |
| Buffer:         | 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2   |  |

### Handling

| Storage:         | 4 °C   |
|------------------|--|
| Storage Comment: | Store lyophilized material at 2-8 °C. For long term storage after reconstitution, prepare small                                |
|                  | aliquots and store at -20 $^{\circ}\text{C}$ . For storage at 2-8 $^{\circ}\text{C}$ , add a preservative to prevent growth of |
|                  | bacteria.  |