

Datasheet for ABIN987823 **Epigen Protein**



Overview Quantity: 25 µg Target: Epigen (EPGN) Origin: Human Escherichia coli (E. coli) Source: **Biological Activity:** Active **Product Details** AVTVTPPITA QQADNIEGPI ALKFSHLCLE DHNSYCINGA CAFHHELEKA ICRCFTGYTG Sequence: **ERCEHLTLTS Y** Characteristics: Determined by the dose-dependent stimulation of the proliferation of murine Balb/3T3 cells. The expected ED50 for this effect is 150-300 ng/ml,corresponding to a specific activity of > 3.3×103 units/mg. > 98 % by SDS-PAGE and HPLC analyses. Purity: Endotoxin Level: Level Less than 1EU/µg of rHuEPG as determined by LAL method **Target Details** Target: Epigen (EPGN) Alternative Name Epigen (EPG) (EPGN Products) Background: Epigen is an EGF-related polypeptide growth factor that signals through the ErbB receptor-1. It is produced in several tissues, including the testis, liver, heart and in certain tumor cells. Epigen

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN987823 | 01/02/2025 | Copyright antibodies-online. All rights reserved.

is mitogenic for fibroblasts and epithelial cells. Human Epigen is initially synthesized as a

Target Details	
	glycosylated 14.7 kDa transmembrane precursor protein, which is processed by proteolytic cleavage to produce a mature soluble sequence. Synonym: Epigen (EPG), Human. Formulation: Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4.
Molecular Weight:	Approximately 7.9kDa monomeric protein, containing 72 amino acid residues, which comprises the EGF homologous portion of the Epigen precursor.
Pathways:	RTK Signaling, EGFR Signaling Pathway
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at < -20 °C. Further dilutions should be made in appropriate buffered solutions.
Storage:	4 °C