## .-online.com antibodies

## Datasheet for ABIN987840 CCL21 Protein



Overview	
Quantity:	1 mg
Target:	CCL21
Origin:	Mouse
Source:	Escherichia coli (E. coli)
Biological Activity:	Active
Product Details	
Sequence:	SDGGGQDCCL KYSQKKIPYS IVRGYRKQEP SLGCPIPAIL FLPRKHSKPE LCANPEEGWV QNLMRRLDQP PAPGKQSPGC RKNRGTSKSG KKGKGSKGCK RTEQTQPSR
Characteristics:	Fully biologically active when compared to standard. The ED50 determined by a chemotaxis bioassay using human peripheral blood lymphocytes is less than 200 ng/ml, corresponding to a specific activity of >, 5.0 × 103 IU/mg.
Purity:	> 97 % by SDS-PAGE and HPLC analyses.
Endotoxin Level:	Level Less than 1EU/ $\mu$ g of rMuExodus-2/CCL21 as determined by LAL method
Target Details	
Target:	CCL21
Alternative Name:	Exodus-2/CCL21 (CCL21 Products)
Background:	Exodus-2/CCL21 is a novel CC chemokine discovered independently by three groups from the EST database, and shows 21-33% identity to other CC chemokines. Exodus-2 contains the four conserved cysteines characteristic of beta chemokines plus two additional cysteines in its

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN987840 | 09/12/2023 | Copyright antibodies-online. All rights reserved.

	unusually long carboxylterminal domain. It is expressed in lymph nodes of certain endothelial cells, and in the spleen and appendix. Exodus-2 chemoattracts T and B lymphocytes and inhibits hematopoiesis. Synonym: Exodus-2/CCL21, Mouse. Formulation: Lyophilized from a 0.2µm filtered concentrated solution in 20mM PB, pH 7.4, 150mM NaCl.
Molecular Weight:	12.0 kDa, a single, non-glycosylated polypeptide chain containing 110 amino acids.
Pathways:	Regulation of Actin Filament Polymerization
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml. Stock solutions should be apportioned into working aliquots and stored at < -20 °C. Further dilutions should be made in appropriate buffered solutions.
Storage:	4 °C