

Datasheet for ABIN987841

CCL21 Protein



_					
	1//	r	Vİ	\triangle	۸/
	V		VI		/ V

Quantity:	20 μg	
Target:	CCL21	
Origin:	Mouse	
Source:	Escherichia coli (E. coli)	
Biological Activity:	Active	
Product Details		
Sequence:	SDGGGQDCCL KYSQKKIPYS IVRGYRKQEP SLGCPIPAIL FLPRKHSKPE LCANPEEGWV	
	QNLMRRLDQP PAPGKQSPGC RKNRGTSKSG KKGKGSKGCK RTEQTQPSR	
Characteristics:	Fully biologically active when compared to standard. The ED50 determined by a chemotaxis	
	bioassay using human peripheral blood lymphocytes is less than 200 ng/ml, corresponding to a	
	specific activity of $>$, 5.0 \times 103 IU/mg.	
Purity:	> 97 % by SDS-PAGE and HPLC analyses.	
Endotoxin Level:	Level Less than 1EU/μg of rMuExodus-2/CCL21 as determined by LAL method	
Target Details		
Target:	CCL21	
Alternative Name:	Exodus-2/CCL21 (CCL21 Products)	
Background:	Exodus-2/CCL21 is a novel CC chemokine discovered independently by three groups from the	
	EST database, and shows 21-33% identity to other CC chemokines. Exodus-2 contains the four	
	conserved cysteines characteristic of beta chemokines plus two additional cysteines in its	

Target Details

- arget betails			
	unusually long carboxylterminal domain. It is expressed in lymph nodes of certain endothelial cells, and in the spleen and appendix. Exodus-2 chemoattracts T and B lymphocytes and inhibits hematopoiesis. Synonym: Exodus-2/CCL21, Mouse. Formulation: Lyophilized from a 0.2µm filtered concentrated solution in 20mM PB, pH 7.4, 150mM NaCl.		
Molecular Weight:	12.0 kDa, a single, non-glycosylated polypeptide chain containing 110 amino acids.		
Pathways:	Regulation of Actin Filament Polymerization		
Application Details			
Restrictions:	For Research Use only		
Handling			
Format:	Lyophilized		
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to t		

bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml. Stock solutions should be apportioned into working aliquots and stored at < -20 °C. Further dilutions should be made in appropriate buffered solutions.

Storage:

4°C