

Datasheet for ABIN988138 **CCL13 Protein**



[Go to Product page](#)

Overview

Quantity:	20 µg
Target:	CCL13
Origin:	Human
Source:	Escherichia coli (E. coli)
Biological Activity:	Active

Product Details

Sequence:	QPDALNVPST CCFTFSSKKI SLQRLKSYVI TTSRCPQKAV IFRTKLGKEI CADPKEKWVQ NYMKHLGRKA HTLK
Characteristics:	Fully biologically active when compared to standard. The ED50 determined by a chemotaxis bioassay using human monocytes is less than 100 ng/ml, corresponding to a specific activity of $>, 1.0 \times 10^4$ IU/mg.
Purity:	> 96 % by SDS-PAGE and HPLC analyses.
Endotoxin Level:	Level Less than 1EU/µg of rHuMCP-4/CCL13 as determined by LAL method

Target Details

Target:	CCL13
Alternative Name:	MCP-4/CCL13 (CCL13 Products)
Background:	CCL13 is a chemoattractant for monocytes and eosinophils, and activates basophils. In addition, it has been reported to be chemotactic for CD4+ and CD8+ T cells, with an activity almost equivalent to that of MCP-3. The bioactivities of CCL13 is most likely mediated by the

Target Details

CC chemokine receptors CCR-2 and CCR-3, both of which have been shown to bind CCL13.
Synonym: MCP-4/CCL13, Human. Formulation: Lyophilized from a 0.2µm filtered concentrated solution in 20mM PB, pH 7.4, 130mM NaCl.

Molecular Weight: 8.6 kDa, a single non-glycosylated polypeptide chain containing 75 amino acids.

Pathways: [Regulation of Systemic Arterial Blood Pressure by Hormones](#), [The Global Phosphorylation Landscape of SARS-CoV-2 Infection](#)

Application Details

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at < -20 °C. Further dilutions should be made in appropriate buffered solutions.

Storage: 4 °C
