

## Datasheet for ABIN988169 **CCL15 Protein**



[Go to Product page](#)

### Overview

Quantity:	25 µg
Target:	CCL15
Origin:	Human
Source:	Escherichia coli (E. coli)
Biological Activity:	Active

### Product Details

Sequence:	QFTNDAETEL MMSKLPLENP VVLNSFHFAA DCCTSYISQS IPCSLMKSYF ETSSECSKPG VIFLTKKGRQ VCAKPSGPGV QDCMKKLPY S
Characteristics:	Fully biologically active when compared to standard. The ED50 determined by a chemotaxis bioassay using human T-lymphocytes is less than 10 ng/ml, corresponding to a specific activity of $>, 1.0 \times 10^5$ IU/mg.
Purity:	$> 97$ % by SDS-PAGE and HPLC analyses.
Endotoxin Level:	Level Less than 1EU/µg of rHuMIP-5/CCL15 as determined by LAL method

### Target Details

Target:	CCL15
Alternative Name:	MIP-5/CCL15 ( <a href="#">CCL15 Products</a> )
Background:	CCL15, a new human CC chemokine, was isolated from a human fetal spleen cDNA library. CCL15 cDNA encodes a predicted 113 amino acid (aa) protein containing a putative signal peptide of 21 amino acids that is cleaved to generate a 92 aa residue mature protein. Within the

## Target Details

---

CC family members, human CCL15 shares 45%, 44%, 35%, and 30% aa homology with mouse C10, human MIPF-1, human HCC-1, and mouse MIP-1<sup>β</sup>, respectively. The gene for MIP-5 is found on chromosome 17 where the genes for most of the human CC chemokines are located. Human CCL15 is expressed in T and B lymphocytes, NK cells, monocytes and monocyte-derived dendritic cells. Human MIP-5 is chemotactic for T cells and monocytes and has been shown to induce calcium flux in human CCR-1-transfected cells. Synonym: MIP-5/CCL15, Human. Formulation: Lyophilized from a 0.2µm filtered concentrated solution in 20mM PB, pH 7.4, 100mM NaCl.

---

Molecular Weight: 10.1 kDa, a single non-glycosylated polypeptide chain containing 92 amino acids.

## Application Details

---

Restrictions: For Research Use only

## Handling

---

Format: Lyophilized

Reconstitution: We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at < -20 °C. Further dilutions should be made in appropriate buffered solutions.

---

Storage: 4 °C