

## Datasheet for ABIN997028

## **HCV Ab ELISA Kit**



_					
	W	0	rv	10	W

Quantity:	96 tests	
Target:	HCV Ab	
Reactivity:	Hepatitis C Virus (HCV)	
Method Type:	Competition ELISA	
Application:	ELISA	
Product Details		
Purpose:	This HCV Ab ELISA test is an enzyme-linked immunosorbent assay for in vitro qualitative	
	identification of IgG antibodies to hepatitis C virus in human serum/plasma.	
Analytical Method:	Qualitative	
Detection Method:	Colorimetric	
Specificity:	96%	
Sensitivity:	99.55%	
Target Details		
Target:	HCV Ab	
Alternative Name:	HCV Ab (HCV Ab Products)	
Target Type:	Antibody, Antibody	
Background:	Hepatitis C virus (HCV) is an envelope, single stranded positive sense RNA (9.5 kb) virus	
	belonging to the family of Flaviviridae. Six major genotypes and series of subtypes of HCV have	
	been identified. Isolated in 1989, HCV is now recognized as the major cause for transfusion	

associated non-A, non-B hepatitis. The disease is characterized with acute and chronic form although more than 50% of the infected individuals develop severe, life threatening chronic hepatitis with liver cirrhosis and hepatocellular carcinomas. Since the introduction in 1990 of anti-HCV screening of blood donations, the incidence of this infection in transfusion recipients has been significantly reduced.

The first generation of HCV ELISAs showed limited sensitivity and specificity and was produced using recombinant proteins complementary to the NS4 (c100-3) region of the HCV genome as antigens. Second generation tests, which included recombinant / synthetic antigens from the Core (c22) and nonstructural regions NS3 (c33c, c100-3) and NS4 (c100-3, c200) resulted in a remarked improvement in sensitivity and specificity. Clinical studies show that significant amount of HCV infected individuals develop antibodies to NS5 non-structural protein of the virus. For this, the third generation tests include antigens from the NS5 region of the viral genome in addition to NS3 (c200), NS4 (c200) and the Core (c22). Third generation tests have improved sensitivity and shorten the time between infection with HCV and the appearance of detectable antibodies (window period) to 60 days.

## Application Details

Sample Volume:	10 μL
Assay Time:	1 - 2 h
Plate:	Pre-coated
Restrictions:	For Research Use only
Handling	
Storage:	4 °C
Expiry Date:	12-14 months